



Risk factors and outcomes for airway failure versus non-airway failure in the intensive care unit: a multicenter observational study of 1514 extubation procedures

Submitted by Beatrice Guillaumat on Mon, 11/19/2018 - 12:58

Titre	Risk factors and outcomes for airway failure versus non-airway failure in the intensive care unit: a multicenter observational study of 1514 extubation procedures
Type de publication	Article de revue
Auteur	Jaber, Samir [1], Quintard, Hervé [2], Cinotti, Raphaël [3], Asehnoune, Karim [4], Arnal, Jean-Michel [5], Guitton, Christophe [6], Paugam-Burtz, Catherine [7], Abback, Paër-Sélim [8], Mekontso Dessap, Armand [9], Lakhali, Karim [10], Lasocki, Sigismond [11], Plantefeve, Gaetan [12], Claud, Bernard [13], Pottecher, Julien [14], Corne, Philippe [15], Ichai, Carole [16], Hajje, Zied [17], Molinari, Nicolas [18], Chanques, Gerald [19], Papazian, Laurent [20], Azoulay, Elie [21], De Jong, Audrey [22]
Editeur	Bohn Stafleu van Loghum
Type	Article scientifique dans une revue à comité de lecture
Année	2018
Langue	Anglais
Date	23 Sept. 2018
Pagination	236
Volume	22
Titre de la revue	Critical Care
ISSN	1574-4280
Mots-clés	Airway [23], Extubation [24], Non-airway [25], weaning [26]

Background

Patients liberated from invasive mechanical ventilation are at risk of extubation failure, including inability to breathe without a tracheal tube (airway failure) or without mechanical ventilation (non-airway failure). We sought to identify respective risk factors for airway failure and non-airway failure following extubation.

Methods

The primary endpoint of this prospective, observational, multicenter study in 26 intensive care units was extubation failure, defined as need for reintubation within 48 h following extubation. A multinomial logistic regression model was used to identify risk factors for airway failure and non-airway failure.

Results

Between 1 December 2013 and 1 May 2015, 1514 patients undergoing extubation were enrolled. The extubation-failure rate was 10.4% (157/1514), including 70/157 (45%) airway failures, 78/157 (50%) non-airway failures, and 9/157 (5%) mixed airway and non-airway failures. By multivariable analysis, risk factors for extubation failure were either common to airway failure and non-airway failure: intubation for coma (OR 4.979 (2.797–8.864), $P < 0.0001$ and OR 2.067 (1.217–3.510), $P = 0.003$, respectively, intubation for acute respiratory failure (OR 3.395 (1.877–6.138), $P < 0.0001$ and OR 2.067 (1.217–3.510), $P = 0.007$, respectively, absence of strong cough (OR 1.876 (1.047–3.362), $P = 0.03$ and OR 3.240 (1.786–5.879), $P = 0.0001$, respectively, or specific to each specific mechanism: female gender (OR 2.024 (1.187–3.450), $P = 0.01$), length of ventilation > 8 days (OR 1.956 (1.087–3.518), $P = 0.025$), copious secretions (OR 4.066 (2.268–7.292), $P < 0.0001$) were specific to airway failure, whereas non-obese status (OR 2.153 (1.052–4.408), $P = 0.036$) and sequential organ failure assessment (SOFA) score ≥ 8 (OR 1.848 (1.100–3.105), $P = 0.02$) were specific to non-airway failure. Both airway failure and non-airway failure were associated with ICU mortality (20% and 22%, respectively, as compared to 6% in patients with extubation success, $P < 0.0001$).

Conclusions

Specific risk factors have been identified, allowing us to distinguish between risk of airway failure and non-airway failure. The two conditions will be managed differently, both for prevention and curative strategies.

Résumé en anglais

URL de la notice

<http://okina.univ-angers.fr/publications/ua18088> [27]

DOI

10.1186/s13054-018-2150-6 [28]

Lien vers le document

<https://ccforum.biomedcentral.com/articles/10.1186/s13054-018-2150-6> [29]

Titre abrégé Crit Care

Liens

- [1] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30455>
- [2] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30456>
- [3] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=22616>
- [4] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=16670>
- [5] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30457>
- [6] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=4907>
- [7] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30458>
- [8] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=31932>
- [9] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30460>
- [10] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30461>
- [11] <http://okina.univ-angers.fr/s.lasocki/publications>
- [12] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30462>

- [13] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30463>
- [14] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30464>
- [15] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30465>
- [16] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30466>
- [17] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30467>
- [18] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30468>
- [19] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30469>
- [20] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=19472>
- [21] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=16340>
- [22] <http://okina.univ-angers.fr/publications?f%5Bauthor%5D=30470>
- [23] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=26046>
- [24] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=26047>
- [25] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=26048>
- [26] <http://okina.univ-angers.fr/publications?f%5Bkeyword%5D=26049>
- [27] <http://okina.univ-angers.fr/publications/ua18088>
- [28] <http://dx.doi.org/10.1186/s13054-018-2150-6>
- [29] <https://ccforum.biomedcentral.com/articles/10.1186/s13054-018-2150-6>

Publié sur *Okina* (<http://okina.univ-angers.fr>)